

A note on the Petri loci

A. BRUNO – E. SERNESI*

Abstract

Let $\mathcal{M}_g$ be the coarse moduli space of complex projective nonsingular curves of genus g . We prove that when the Brill-Noether number $\rho(g, r, n)$ is non-negative every component of the Petri locus $P_{g,n}^r \subset \mathcal{M}_g$ whose general member is a curve C such that $W_n^{r+1}(C) = \emptyset$, has codimension one in $\mathcal{M}_g$.

1 Introduction

Let C be a nonsingular irreducible projective curve of genus $g \geq 2$ defined over $\mathbb{C}$. A pair (L, V) consisting of an invertible sheaf L on C and of an $(r+1)$ -dimensional vector subspace $V \subset H^0(L)$, $r \geq 0$, is called a *linear series* of dimension r and degree $n = \deg(L)$, or a g_n^r . If $V = H^0(L)$ then the g_n^r is said to be *complete*.

If (L, V) is a g_n^r then the *Petri map* for (L, V) is the natural multiplication map

$$\mu_0(L, V) : V \otimes H^0(\omega_C L^{-1}) \longrightarrow H^0(\omega_C)$$

The Petri map for L is

$$\mu_0(L) : H^0(L) \otimes H^0(\omega_C L^{-1}) \longrightarrow H^0(\omega_C)$$

Recall that C is called a *Petri curve* if the Petri map $\mu_0(L)$ is injective for every invertible sheaf L on C . By the Gieseker-Petri theorem [5] we know that in $\mathcal{M}_g$, the coarse moduli space of nonsingular projective curves of genus g , the locus of curves which are not Petri is a proper closed subset P_g , called the *Petri locus*. This locus decomposes into several components, according to the numerical types and to other properties that linear series can have on a curve of genus g . We will say that C is *Petri with respect to g_n^r 's* if the Petri map $\mu_0(L, V)$ is injective for every $g_n^r (L, V)$ on C .

We denote by $P_{g,n}^r \subset \mathcal{M}_g$ the locus of curves which are not Petri w.r. to g_n^r 's. Then

$$P_g = \bigcup_{r,n} P_{g,n}^r$$

*Both authors are members of GNSAGA-INDAM

where the union is finite by obvious reasons. The structure of $P_{g,n}^r$ and of P_g is not known in general: both might a priori have several components and not be equidimensional. In some special cases $P_{g,n}^r$ is known to be of pure codimension one (notably in the obvious case $\rho(g, r, n) = 0$, and for $r = 1$ and $n = g - 1$ [12]). If the Brill-Noether number

$$\rho(g, r, n) := g - (r + 1)(g - n + r)$$

is nonnegative then it is natural to conjecture that $P_{g,n}^r$ has pure codimension one if it is non-empty. The evidence is the fact that $P_{g,n}^r$ is the image in $\mathcal{M}_g$ of a determinantal scheme $\tilde{P}_{g,n}^r$ inside the relative Brill-Noether scheme $\mathcal{W}_n^r \rightarrow \mathcal{M}_g$, and the expected dimension of $\tilde{P}_{g,n}^r$ is $3g - 4$. This is the point of view that we apply for the proof of our main theorem 1.1 (see below). One might ask if even P_g has pure codimension one: there is not much evidence for this, except that it can be directly checked to be true for low values of g (see the very recent preprint by M. Lelli-Chiesa [10]).

Before stating our result we recall what is known. Denote by $\overline{\mathcal{M}}_g$ the moduli space of stable curves, and let

$$\overline{\mathcal{M}}_g \setminus \mathcal{M}_g = \Delta_0 \cup \cdots \cup \Delta_{[\frac{g}{2}]}$$

be its boundary, in standard notation. In [2] G. Farkas has proved the existence of at least one divisorial component of $P_{g,n}^1$ in case $\rho(g, 1, n) \geq 0$ and $n \leq g - 1$, using the theory of limit linear series. He found a divisorial component which has a nonempty intersection with Δ_1 . Another proof has been given in [1], by degeneration to a stable curve with g elliptic tails. The method of [2] has been extended in [3] to arbitrary r . In this note without using any degeneration argument we prove the following result:

Theorem 1.1 *If $\rho(g, r, n) \geq 0$ then every component of $P_{g,n}^r$ whose general member is a curve C such that $W_n^{r+1}(C) = \emptyset$, has codimension one in $\mathcal{M}_g$.*

Note that a necessary numerical condition for the existence of a curve C as in the statement is that $\rho(g, r + 1, n) < 0$. This condition, together with $\rho(g, r, n) \geq 0$ gives:

$$0 \leq \rho(g, r, n) < g - n + 2(r + 1)$$

or, equivalently:

$$\frac{r}{r+1}g + r \leq n < \frac{r+1}{r+2}g + r + 1$$

For the proof of the theorem we introduce a modular family $\mathcal{C} \rightarrow B$ of curves of genus g (see (i) below for the definition) and we use the determinantal description of the relative locus $\mathcal{W}_n^r(\mathcal{C}/B)$ over B and of the naturally defined closed subscheme $\tilde{P}_{g,n}^r \subset \mathcal{W}_n^r(\mathcal{C}/B)$ whose image in $\mathcal{M}_g$ is $P_{g,n}^r$. Since it is a determinantal locus, every component of $\tilde{P}_{g,n}^r$ has dimension $\geq 3g - 4$. Then a theorem of F. Steffen [11] ensures that every component of $P_{g,n}^r$ has dimension $\geq 3g - 4$ as well, thus proving the result.

In a forthcoming paper (in preparation) we will show the existence of a divisorial component of $P_{g,n}^1$ which has a non-empty intersection with Δ_0 , when $\rho(g, 1, n) \geq 1$.

2 Proof of Theorem 1.1

In this section we fix g, r, n such that $\rho(g, r, n) \geq 0$ and $\rho(g, r + 1, n) < 0$. Consider the following diagram:

$$\begin{array}{ccc} J_n(\mathcal{C}/B) \times_B \mathcal{C} & \longrightarrow & \mathcal{C} \\ \downarrow & & \downarrow f \\ J_n(\mathcal{C}/B) & \xrightarrow{q} & B \end{array} \quad (1)$$

where:

- (i) f is a smooth modular family of curves of genus g parametrized by a non-singular quasi-projective algebraic variety B of dimension $3g - 3$. This means that at each closed point $b \in B$ the Kodaira-Spencer map $\kappa_b : T_b B \rightarrow H^1(\mathcal{C}(b), T_{\mathcal{C}(b)})$ is an isomorphism. In particular, the functorial morphism

$$\beta : B \longrightarrow \mathcal{M}_g$$

is finite and dominant. The existence of f is a standard fact, see e.g. [7], Theorem 27.2.

- (ii) $J_n(\mathcal{C}/B)$ is the relative Picard variety parametrizing invertible sheaves of degree n on the fibres of f .
- (iii) For all closed points $b \in B$ the fibre $\mathcal{C}(b)$ satisfies $W_n^{r+1}(\mathcal{C}(b)) = \emptyset$. This condition can be satisfied modulo replacing B by an open subset if necessary, because the condition $W_n^{r+1}(\mathcal{C}(b)) = \emptyset$ is open w.r. to $b \in B$.
- (iv) We may even assume that any given specific curve C of genus g satisfying $W_n^{r+1}(C) = \emptyset$ appears among the fibres of f . In particular we may assume that the dense subset $\text{Im}(\beta) \subset \mathcal{M}_g$ has a non-empty intersection with all irreducible components of $P_{g,n}^r$ whose general element parametrizes a curve C such that $W_n^{r+1}(C) = \emptyset$.

Let $\mathcal{P}$ be a Poincaré invertible sheaf on $J_n(\mathcal{C}/B) \times_B \mathcal{C}$. Using $\mathcal{P}$ in a well-known way one constructs the relative Brill-Noether scheme

$$\mathcal{W}_n^r(\mathcal{C}/B) \subset J_n(\mathcal{C}/B)$$

Consider the restriction of diagram (1) over $\mathcal{W}_n^r(\mathcal{C}/B)$:

$$\begin{array}{ccc} \mathcal{W}_n^r(\mathcal{C}/B) \times_B \mathcal{C} & \xrightarrow{p_2} & \mathcal{C} \\ p_1 \downarrow & & \downarrow f \\ \mathcal{W}_n^r(\mathcal{C}/B) & \xrightarrow{q} & B \end{array} \quad (2)$$

Every irreducible component of $\mathcal{W}_n^r(\mathcal{C}/B)$ has dimension $\geq 3g-3+\rho(g, r, n)$ and, since $\rho(g, r, n) \geq 0$, there is a component which dominates B [8, 9]. A closed point $w \in \mathcal{W}_n^r(\mathcal{C}/B)$ represents an invertible sheaf L_w on the curve $\mathcal{C}(q(w))$ such that $h^0(L_w) \geq r+1$. Denoting again by $\mathcal{P}$ the restriction of $\mathcal{P}$ to $\mathcal{W}_n^r(\mathcal{C}/B) \times_B \mathcal{C}$, we have a homomorphism of coherent sheaves on $\mathcal{W}_n^r(\mathcal{C}/B)$, induced by multiplication of sections along the fibres of p_1 :

$$\mu_0(\mathcal{P}) : p_{1*}\mathcal{P} \otimes p_{1*}[p_2^*(\omega_{\mathcal{C}/B}) \otimes \mathcal{P}^{-1}] \longrightarrow p_{1*}[p_2^*\omega_{\mathcal{C}/B}]$$

By condition (iii) above, these sheaves are locally free, of ranks $(r+1)(g-n+r)$ and g respectively. Moreover, by definition, at each point $w \in \mathcal{W}_n^r(\mathcal{C}/B)$, the map $\mu_0(\mathcal{P})$ coincides with the Petri map

$$\mu_0(L_w) : H^0(\mathcal{C}(q(w)), L_w) \otimes H^0(\mathcal{C}(q(w)), \omega_{\mathcal{C}(q(w))} L_w^{-1}) \longrightarrow H^0(\mathcal{C}(q(w)), \omega_{\mathcal{C}(q(w))})$$

Claim: the vector bundle

$$[p_{1*}\mathcal{P} \otimes p_{1*}[p_2^*(\omega_{\mathcal{C}/B}) \otimes \mathcal{P}^{-1}]]^\vee \otimes p_{1*}[p_2^*\omega_{\mathcal{C}/B}] \quad (3)$$

is q -relatively ample.

Proof of the Claim. If we restrict diagram (2) over any $b \in B$ and we let $C = \mathcal{C}(b)$, we obtain:

$$\begin{array}{ccc} W_n^r(C) \times C & \xrightarrow{\pi_2} & C \\ \pi_1 \downarrow & & \\ W_n^r(C) & & \end{array}$$

and the map $\mu_0(\mathcal{P})$ restricts over $W_n^r(C)$ to

$$m_P : \pi_{1*}P \otimes \pi_{1*}[\pi_2^*\omega_C \otimes P^{-1}] \longrightarrow H^0(C, \omega_C) \otimes \mathcal{O}_{W_n^r(C)}$$

where $P = \mathcal{P}|_{W_n^r(C) \times C}$ is a Poincaré sheaf on $W_n^r(C) \times C$. The dual of the source of m_P is an ample vector bundle (compare [4], §2), while the target is a trivial vector bundle, and therefore

$$[\pi_{1*}P \otimes \pi_{1*}[\pi_2^*\omega_C \otimes P^{-1}]]^\vee \otimes_{\mathbb{C}} H^0(C, \omega_C)$$

is an ample vector bundle. This means that the vector bundle (3) restricts to an ample vector bundle on the fibres of q . This implies, by [6], Th. 4.7.1, applied

to the invertible sheaf $\mathcal{O}(1)$ on the projective bundle associated to (3), that (3) is q -relatively ample. This proves the Claim.

Consider the degeneracy scheme:

$$\tilde{P}_{g,n}^r := D_{(r+1)(g-n+r)-1}(\mu_0(\mathcal{P})) \subset \mathcal{W}_n^r(\mathcal{C}/B)$$

which is supported on the locus of $w \in \mathcal{W}_n^r(\mathcal{C}/B)$ such that $\mu_0(L_w)$ is not injective. By applying Theorem 0.3 of [11] to it we deduce that every irreducible component of $q(\tilde{P}_{g,n}^r) \subset B$ has dimension at least

$$\dim[\mathcal{W}_n^r(\mathcal{C}/B)] - [g - (r+1)(g-n+r) + 1] = 3g - 4$$

Since f is a modular family, it follows that every irreducible component of $\beta(q(\tilde{P}_{g,n}^r)) \subset \mathcal{M}_g$ has dimension $\geq 3g - 4$ as well. But $\beta(q(\tilde{P}_{g,n}^r)) \subset P_{g,n}^r \neq \mathcal{M}_g$ and therefore all the components of $\beta(q(\tilde{P}_{g,n}^r))$ are divisorial. Since, by (iv), $\beta(q(\tilde{P}_{g,n}^r))$ is the union of all the components of $P_{g,n}^r$ whose general element parametrizes a curve C such that $W_n^{r+1}(C) = \emptyset$, the theorem is proved. $\square$

References

- [1] A. Castorena, M. Teixidor i Bigas: Divisorial components of the Petri locus for pencils, *J. Pure Appl. Algebra* 212 (2008), 1500–1508.
- [2] G. Farkas: Gaussian maps, Gieseker-Petri loci and large theta-characteristics, *J. reine angew. Mathematik* 581 (2005), 151-173.
- [3] G. Farkas: Rational maps between moduli spaces of curves and Gieseker-Petri divisors, *Journal of Algebraic Geometry* 19 (2010), 243-284.
- [4] W. Fulton - R. Lazarsfeld: On the connectedness of degeneracy loci and special divisors, *Acta Math.* 146 (1981), 271-283.
- [5] D. Gieseker: Stable curves and special divisors, *Inventiones Math.* 66 (1982), 251-275.
- [6] A. Grothendieck: *Elements de Geometrie Algebrique* III, 1. Inst. de Hautes Etudes Sci. Publ. Math. 11, 1961.
- [7] R. Hartshorne: *Deformation Theory*, Springer GTM vol.257 (2010).
- [8] Kempf G.: Schubert methods with an application to algebraic curves, *Publication of Mathematisch Centrum*, Amsterdam 1972.
- [9] Kleiman S., Laksov D.: On the existence of special divisors, *Amer. Math. J.* 94 (1972), 431-436.
- [10] M. Lelli-Chiesa: The Gieseker-Petri divisor in M_g for $g \leq 13$. *arXiv* preprint n. 1012.3061v1.

- [11] F. Steffen: A generalized principal ideal theorem with an application to Brill-Noether theory. *Inventiones Math.* 132 (1998), 73-89.
- [12] M. Teixidor: The divisor of curves with a vanishing theta null, *Compositio Math.* 66 (1988), 15-22.

ADDRESS OF THE AUTHORS:

Dipartimento di Matematica, Università Roma Tre
Largo S. L. Murialdo 1, 00146 Roma, Italy.

`bruno@mat.uniroma3.it`

`sernesi@mat.uniroma3.it`